

RESEARCH ARTICLE

Steroid injections in musculoskeletal conditions and COVID infection rates: What is the impact on positive rates following the injection?

Daniel Lethbridge^{1,2}  | Simon O'Shea³¹Healthshare Ltd, Kent, West Malling, UK²Physiotherapy Department, University of Hull, Hull, UK³Physiotherapy Department, Sheffield Hallam University, Sheffield, UK**Correspondence**

Daniel Lethbridge, Physiotherapy Department, University of Hull, Cottingham Road, Hull, HU67RX, UK.
Email: d.lethbridge@hull.ac.uk

Abstract

Background and Aim: Therapeutic glucocorticosteroid injections are commonly utilised to manage musculoskeletal (MSK) complaints. Following the SARS-CoV-2 pandemic, national guidelines advised against their use due to potential immunosuppressant effects. The aim of the study was to determine whether steroid injections for MSK conditions impacts on positive COVID 19 infection rates.

Patients and Methods: This retrospective evaluation involved primary care participants who received a steroid injection for a MSK condition. 291 participants receiving a total of 299 steroid injections entered the study between the 25 September 2020 and the 29 April 2021.

Results: Six participants had positive polymerase chain reaction (PCR) tests, averaging 22.83 days (SD 10.48) after the injection. An infection rate of 2.06% was demonstrated in the injection group with the control group demonstrating 6.97% ($p = 0.000752$) with statistical significance set at $p = 0.05$. The odds ratio was identified as 0.27 indicating a lower odds of a positive PCR test compared with the control group.

Conclusions: This retrospective evaluation found a low risk of positive PCR tests for low and moderate COVID-19 risk patients injected during the COVID-19 pandemic. Glucocorticosteroid injections within the COVID-19 pandemic were not associated with higher COVID-19 rates compared to the local population, in fact, they were related to lower rates. For future studies, large scale studies and meta analyses are needed to provide greater generalisation to the population.

KEYWORDS

2019-ncov, coronavirus, corticosteroid, COVID-19, glucocorticosteroid

1 | INTRODUCTION

In December 2019 an atypical severe acute respiratory syndrome occurred within Wuhan, China and rapidly spread throughout the world creating a Worldwide Pandemic. The disease referred to as COVID-19 was caused by a novel coronavirus known as SARS-CoV-2. The SARS-CoV-2 infection can trigger both innate and adaptive

human immune system responses which if uncontrolled can lead to local and systemic tissue damage (Cao, 2020). The virus can activate immune responses and trigger significant antibody production along with significantly elevating pro-inflammatory cytokines levels (Cao, 2020). Various symptoms are recognised in response to COVID-19 infection and these are predominantly respiratory system symptoms but can affect multiorgan systems and have led to

significant fatalities (Yuki et al., 2020). In January 2020 the first positive COVID-19 case was found in the United Kingdom causing significant impact on a wide spectrum of services, none the least the NHS. By March 2020 a National Lockdown was introduced and people were told to work from home and not travel unless absolutely necessary. As a result, face to face clinics within musculoskeletal (MSK) Physiotherapy services including injection clinics, were ceased.

Musculoskeletal condition management in Physiotherapy, Orthopaedic and Rheumatological services utilise glucocorticosteroid injections (CSIs) as adjuncts to help manage multiple MSK and inflammatory disorders. Glucocorticosteroids play vital roles in maintaining homeostasis and influencing innate immune responses and are recognised as providing both anti-inflammatory and pro-inflammatory responses through influencing glucocorticoid receptors via various mechanisms (Cruz-Topete & Cidlowski, 2015). They provide their anti-inflammatory effect by reducing inflammatory mediator production and release, causing vascular reaction suppression that occurs during the inflammatory response (Becker, 2013). The immunosuppressant effects predominantly occur through their effect on the hypothalamic-pituitary adrenal axis (HPA) and their inhibitory effects on macrophages and T cells as well as leucocyte function (Becker, 2013). Inflammation is common within MSK conditions therefore glucocorticosteroids are potent anti-inflammatories that are routinely administered in MSK practice to reduce inflammation, manage pain and improve function (Stephens et al., 2008).

Following the pandemic outbreak various societies (British Pain Society, 2020; British Society of Rheumatology, 2020; British Society of Skeletal Radiology, 2020; Faculty of Pain Medicine of the Royal College of Anaesthetists, 2020) produced National guidelines regarding CSI use due to their immunosuppressant effects and subsequent potential increased risk of developing COVID-19. These guidelines clearly highlight that injections should only be considered with severe disease activity or with significant levels of pain and disability and in cases with no alternative treatments (BSR, BOA, BASS, RCGP, BSIR, FPM, BPS, CSP, 2020). Further guidance advocated against injection therapy for vulnerable patient groups, that is, those over the age of 70 with co-morbidities such as diabetes, ischaemic heart disease or chronic respiratory disorders as they are classified as high risk of developing COVID-19 (BSR, BOA, BASS, RCGP, BSIR, FPM, BPS, CSP, 2020).

Within the current study the most commonly utilised glucocorticosteroid for managing MSK complaints was Triamcinolone Acetonide and this study utilised this drug solely. Triamcinolone Acetonide has been demonstrated to cause adrenal suppression within the first 48 h following administration (Fascia et al., 2020) with these effects continuing for 30–40 days (Broersen et al., 2015; Fascia et al., 2020). Broersen et al. (2015) found that adrenal suppression with the use of corticosteroids has been particularly apparent following administration of an intra-articular joint injection with 52% of patients developing adrenal insufficiency following joint injection. With the recognised immunosuppressant effects, it remains unclear the potential impact that this suppression may have and whether it

leads to increased coronavirus infection at the time of injection, or within the proceeding 40 days (BSR, BOA, BASS, RCGP, BSIR, FPM, BPS, CSP, 2020).

This evaluation, therefore, aimed to ascertain whether any significant risk was apparent providing evidence for, or against, the use of corticosteroid injections during such pandemic situations, providing useful evidence to aid in shared decision making with regards to administering CSIs.

1.1 | Aims

1. Steroid injections in MSK conditions and COVID infection rates: what is the impact on positive rates following the injection?
2. To determine whether COVID 19 positive results differed between risk category (mild or moderate).
3. To determine whether there was any relationship between steroid dose and those demonstrating a positive polymerase chain reaction (PCR) test for COVID-19 following a CSI.

2 | RESEARCH DESIGN & METHODS

The evaluation utilised data routinely gathered by a Healthcare organisation. A report containing a list of patients injected within the service between 25 September 2020 and 29 April 2021 was obtained. This timeframe was chosen as the start date was recommencing injection therapy within the Healthcare organisation following the original cessation at the start of the pandemic and the end date was via convenience sampling as it was the latest date available at the time of data collection, which provided the greatest number of potential participants.

Each participant was given a reference number which was recorded along with data on injection date, body part injected, dose and patient demographics.

The control group was obtained through data provided by the Office of National Statistics (2020) which provided daily COVID-19 case numbers for the Hull region within the United Kingdom, for the duration of the data collection period. This provided accurate daily cases and up to date population data that was used to calculate the average number of positive COVID-19 cases within the Hull region. The control and intervention groups were both obtained from the Hull region making the groups more reflective of each other and allowing direct comparison of COVID-19 levels.

A review of the patient records within the General Practice database was performed by the author to obtain records of any positive PCR COVID-19 tests within a 40-day period following corticosteroid injection. Further data regarding the COVID-19 risk group of each patient was obtained and was based around the NHS risk criteria (NHS Digital, 2021). Positive COVID-19 PCR test data was recorded alongside the date of the positive test, the number of days elapsed between the positive test and the injection and the who had a positive COVID-19 test following injection.

2.1 | Participants

National guidelines and company policy during the Pandemic meant that participants were excluded from injection if they were considered high risk of COVID-19 development as per the NHS risk category (NHS Digital, 2021) or were under the age of 18 years. The study utilised low and moderate COVID 19 risk patients indicating that they were systemically well without any significant past medical history (including active inflammatory arthropathies) and were not taking immunosuppressant medication. The vast majority of the subjects were injected into a single joint/soft tissue and other than the presenting MSK condition were otherwise fit and healthy. Retrospective convenience sampling was used to obtain the participants as its emphasis was based on a known population of people injected within the MSK service and could not, therefore, be truly random. The sample size was determined via convenience sampling as it encompassed the greatest number of participants available at the time of data collection.

3 | FINDINGS/RESULTS

Between 25 September 2020 and 29 April 2021 (216 days), 575 triamcinolone steroid injections were administered to a total of 560 patients. Unfortunately, due to a lack of General Practice data sharing, access to full patient records was only available for 291 participants receiving a total of 299 steroid injections. Of the 291 participants 183 were female (62.89%) and 108 were male (37.11%) with an average age of 59.73 years (range 33–88 years, standard deviation [SD] 11.15). Within the injection group, 184 participants (63.2%) were in the low risk of COVID-19 development group and 107 (36.8%) in the moderate risk group.

Various soft tissue and joint injections were administered for multiple MSK complaints during the evaluation period with varying doses of triamcinolone acetonide used with an average dose per injection of 19.44 mg (range 10–40 mg, SD 12.11). The steroids were injected independently or diluted with either lidocaine hydrochloride 1% or 2% or sodium chloride 0.9% in the most frequently injected sites were the shoulder (23.1%) and knee joints (22.1%) making up 45.2% of the total injections with a mean dose of 40 mg of triamcinolone acetonide per injection. Of the 299 injections 150 (50.2%) were performed under ultrasound guidance and 149 (49.8%) were blind injections. The site and injection doses utilised are listed in Table 1.

The control group contained 259,481 participants, which was the regional population and over the study duration 18,088 participants recorded positive PCR tests [12] (absolute risk = 6.97%) of the regional population testing positive for COVID-19.

The General Practice database for each participant who had received a steroid injection was reviewed for positive PCR COVID-19 tests within 40-day of the injection. Six participants (absolute risk = 2.06%) were identified as having a positive PCR test within 40-day with four being male and two being female (Figure 1). The number need to treat was calculated as 20.12.

TABLE 1 Site and dose of triamcinolone acetonide injection

Site of injection	Number of injections	Steroid dose in mg
Acromioclavicular joint	8	10
Carpal tunnel	13	20
Carpometacarpal joint	15	10
De Quervain's tenosynovitis	5	10
Elbow joint	2	10
First metatarsal joint	2	20
Foot injection	1	10
Greater trochanteric pain	8	20–40
Golfers elbow	2	10
Hip joint	17	40
Knee joint	66	40
Lateral coronary ligament knee	1	40
Morton's neuroma	9	20
Plantar fascia injection	12	15–20
Shoulder joint	69	40
Subacromial bursa	39	20
Tarsometatarsal joint	6	20
Tennis elbow	7	10
Trigger finger/thumb	16	10
Wrist joint	1	10
		Mean dose: 19.44
		Range 10–40

The average age of these participants was 56.83 years (range 45–63 years, SD 6.85). Five out of the six participants (83.33%) were in the low COVID-19 risk group and one participant (16.67%) was in the moderate COVID-19 risk group (Figure 1). The injection sites varied in location within this group with two participants (33.33%) receiving shoulder joint injections and the remaining four receiving injections into different soft tissues and joints (Table 2). Ultrasound guidance was used to perform 4 (66.67%) out of the six injections with 2 (33.33%) performed blind. The average dose of triamcinolone acetonide injected was 28.33 mg (range 20–40 mg, SD 9.83) per injection for this group and each injection with the exception of one was diluted with lidocaine hydrochloride 1% or 2% ranging from 1 to 9 ml.

The average time that elapsed between the injection and positive PCR COVID-19 test was 22.83 days (range 7–33 days, SD 10.48). 4 (66.67%) out of 6 participants received their injection when case numbers were relatively low and the remaining 2 (33.33%) received their injection during the peak of positive PCR tests when the infection rate was at its highest within the study period (Figure 2).

The Fisher's exact test demonstrated a p -value of <0.05 ($p = 0.0002$) with an odds ratio of 0.27 (Table 3).




FIGURE 1 Timeline of the number of days passed between glucocorticosteroid injection, positive COVID 19 test and the COVID 19 risk category

TABLE 2 Intervention group positive COVID-19 participant injection sites

Injection site	Number	Dose of steroid	USGI or blind
Subacromial bursa	1	20	USGI
Carpal tunnel	1	20	Blind
Plantar fascia	1	20	Blind
Shoulder joint	2	30/40	USGI (x2)
Hip joint	1	40	USGI

Abbreviation: USGI, ultrasound guided injection.

4 | DISCUSSION

At the start of the COVID-19 pandemic there was significant concern over the potential immunosuppressant effects and subsequent COVID-19 risk posed following corticosteroid therapy. This study demonstrates a low incidence within the UK based cohort of a positive COVID-19 PCR test following a corticosteroid injection for a MSK complaint. There were only 6 positive PCR tests out of 299 injections with an infection rate of 2.06% which when compared with the infection rate of 6.97% positive COVID-19 tests within the control group. The Fisher's exact test demonstrated a p -value of <0.05 ($p = 0.0002$) indicating dependent variables given the statistically significant difference between those injected and the control group. The odds ratio was 0.27, which being <1 indicates that the exposure (CSI) was associated with lower odds of the tested outcome (positive PCR test for COVID-19) when compared with the control group. This highlights that injection of triamcinolone acetonide was a low risk procedure for patients within the low and moderate COVID-19 risk groups and interestingly the infection rate was less than that of the local population. This could be due to the potentially beneficial effects of glucocorticosteroids in treating COVID-19 as Finney et al. (2021) demonstrated. They noted a potential reduction in susceptibility of chronic obstructive pulmonary disease patients receiving inhaled corticosteroids due to a reduction in SARS-CoV-2 entry receptor ACE2 however given the greater dose, regularity

and method of delivery of the inhaled steroid it may be feasible that this reduction in infection rate may well have occurred due to other causes. Extensive counselling and written consent process including the discussion of the risks and benefits of the CSI prior to receiving the injection may have led to greater caution being shown by the participants and taking more precautions with regards to social contact and self-isolation/distancing and carrying out more effective hand hygiene measures in the knowledge that the steroid can reduce the immune response. The findings of this study are consistent with and in keeping with other studies (Aziz et al., 2021; Bugeja et al., 2021; Chang et al., 2021; McKean et al., 2020; Morgan & Dattani, 2020; Newton et al., 2021; Regan et al., 2021). The risk identified within those studies demonstrated extremely low infection risk with very few or no positive cases of COVID-19 infection following CSI.

Since the initiation of the presented study others have also investigated this topic. One example (Bugeja et al., 2021) carried out a retrospective study on 734 participants who received a CSI for a MSK complaint and reviewed any subsequent COVID-19 development. The study had chosen a 30-day timeframe for the action of the steroid which may have limited the number of positive cases highlighted. They found no increased risk of contracting COVID-19 following CSI. The study used an intervention group as well as a matched, randomly selected control group. Appropriate statistical testing using the Fisher's exact test was performed on the data given the relatively low sample size and no statistically significant difference ($p < 0.05$) was apparent between the intervention group and control group. Four participants developed COVID-19 within 30 days with three out of the four receiving injections into more than one body part. This may indicate that they received a higher dose of corticosteroid than those with single joint/soft tissue injections and subsequently at higher risk of HPA axis suppression, as a dose dependent relationship exists with regards to higher doses of glucocorticosteroid leading to greater levels of adrenal suppression (Habib, 2009).

Chang et al. (2021) within their prospective study of 66 patients undergoing image guided corticosteroid injections demonstrated no




FIGURE 2 Date of confirmed COVID-19 cases in North of England Region also indicating date steroid injections were administered for positive COVID-19 participants

TABLE 3 Odds ratios and Fisher's exact test results

	Positive COVID-19 PCR test	Negative or no COVID-19 PCR test	Total	Odds
CSI positive PCR test group	6	293	299	0.02
Control group	18,088	241,393	259,481	0.07
Total	18,094	241,686		
Results	Odds ratio = 0.27		259,780 (grand total)	
	Fisher's exact test $p = 0.0002$			

Abbreviation: PCR, polymerase chain reaction.

statistically significant difference in COVID-19 infection rates when compared with the general population. Unfortunately, only low participant numbers were recruited limiting the statistical power of the results ($n = 66$). The study was also carried out during the second part of the lockdown period whereby the rate of COVID-19 infections within the general population would have been low and at that time widespread testing was not available for the general population indicating that a true COVID-19 infection rate for this particular population was unknown. Although limitations were apparent, only one out of 66 participants (1.52%) went on to develop COVID-19, 19 days following their injection with no other patients reporting symptoms or testing positive for COVID-19.

McKean et al. (2020) within their retrospective observational study of 504 CSIs reported very low incidences of positive COVID-19 infections following a CSI with no adverse clinical outcomes. Of the 504 injections only 11 COVID-19 tests were performed on nine patients with no positive results found reinforcing the low risk of the procedure, however the participant numbers were relatively low limiting the determination of any absolute risk. The study referred to Upper Tier Local Authority (UTLA) COVID-19 infection rates as a comparison however no direct control group was used and the infection rates within the UTLA would have included patients within

high COVID-19 risk groups. These patients were excluded within this study as the national guidance advised against utilising CSIs for this particular patient group due to the elevated COVID-19 development risk. Therefore, the comparison needs to take into account that this particular infection rate may have varied for their population group. Within both the McKean et al. (2020) study and also the present study an element of bias may have been introduced as they were based on specific local populations with unique demographics in terms of socioeconomic group and race. The national guidance on informed consent prior to CSI and the requirement to explain in depth the potential risk may also have influenced the participants leading them to self-isolate and demonstrate greater COVID-19 precautions than those within the general population potentially affecting the risk posed due to the reduced contact with COVID-19.

Since the onset of the COVID-19 pandemic several studies (RECOVERY Collaborative Group, 2021; Yu et al., 2021) have been published relating to the treatment of COVID-19 with oral, inhaled and intravenously administered glucocorticoids. The PRINCIPLE study (Yu et al., 2021) in their randomised controlled trial of 4700 participants utilised inhaled budesonide in the treatment of COVID-19 patients and compared this to usual care and usual care with an alternative treatment and found improvement in time to

recovery in higher risk complication groups receiving this therapy compared to those who did not however the probability to superiority was below the threshold specified. Further support for corticosteroids in the treatment of COVID-19 was found within The RECOVERY trial (RECOVERY Collaborative Group, 2021) in their randomised controlled trial of 9355 participants who compared patients receiving dexamethasone with usual care for hospitalised patients with COVID-19 and found a reduced 28-day mortality rate with oral or intravenous dexamethasone in COVID-19 patients receiving oxygen therapy and mechanical ventilation and this has since been recommended as treatment for these patients via NHS England (2020).

4.1 | Limitations

There are several limitations to this study starting primarily with relatively low participants numbers which unfortunately was compounded by the lack of General Practice data sharing which significantly reduced total participant numbers. There was potential risk of bias given the very specific local participant demographics used in terms of ethnicity and socioeconomic group making it more difficult to generalise the findings to the general population.

This evaluation relied on accurate COVID-19 testing in the form of PCR tests and these have a pre-test probability of 80%, a sensitivity of 70% and a specificity of 95% (Watson et al., 2020) leading to understandable false positives and false negatives in the testing. As well as accepting the lack of total accuracy of the testing, reliance upon patients actually attending for swab testing when they had symptoms was also a potential limitation however it is hoped this was consistent between both groups. Furthermore, not all participants were tested for COVID-19, leaving the potential that some injected participants may have developed COVID-19 but were asymptomatic given that a high proportion ranging from 20% to 75% of COVID-19 infections are asymptomatic (Yanes-Lane et al., 2020).

The control group included participants that tested positive for COVID-19 and may have been in the high risk of developing COVID-19 category and these were excluded from this study, which may have influenced the infection rate within the local population. The data utilised for the control group may also have included positive lateral flow tests as the PCR testing was usually carried out following a positive lateral flow test and, therefore, some of the positive COVID-19 cases may have been awaiting PCR tests. Unfortunately, it is feasible that some of the participants within the intervention group may have had positive tests in different regions within the UK and that these may not have been recorded in their General Practice patient notes, however England had its second and third national lockdowns during the study period with various restrictions placed upon general movement, therefore, it is hoped that any missing data would be negligible. Given the nature of the consent process and guidelines put in place during the pandemic, even greater emphasis was placed on

explicitly detailing the risks of the CSI making very specific reference to the immunosuppressant effects and the subsequent risk of developing COVID-19 with written consent being gained prior to the procedure. Telephone consultations prior to the CSI were carried out informing the patient of the risks and advising self-isolation and social distancing following the CSI which may have resulted in the participants demonstrating greater levels of distancing from society in fear of developing COVID-19. The timeframe utilised within the study was set at 40 days as several studies (Broersen et al., 2015; Fascia et al., 2020) had highlighted a 30–40 days duration of steroid action therefore the highest reported duration of action of the steroid was utilised.

5 | CONCLUSION

This retrospective evaluation found a low risk of a positive PCR test for COVID-19 for participants within low and moderate COVID-19 risk categories, injected with triamcinolone acetonide for a MSK complaint. None of the participants that tested positive for COVID-19 following a CSI had had a vaccination prior to their injection leading to reinforcement that the procedure appears to be low risk. The COVID-19 infection risk within the control group was greater than those receiving a CSI, providing confirmation that CSI for this patient group was low risk. The study demonstrates that there was no greater risk of a positive COVID-19 PCR test with increasing doses of triamcinolone acetonide and there was no evidence of any increased risk whether the injection undertaken was targeted at soft tissue or administered within local joints. A very small increased rate of positive PCR tests were associated with those injected under ultrasound guidance, however, the participant numbers were too low to draw any significant conclusions. There was no evidence that increasing age increased the risk of positive PCR tests following CSI or that participants within the moderate risk of developing COVID-19 categories were at greater risk of positive PCR results. There was a slight increased rate of a positive PCR test in males with a greater number of participants testing positive following a CSI but the low numbers prevent inference about gender differences.

The findings of this study add to the current base of literature concerning the low risk of utilising CSI injection therapy during the COVID-19 pandemic. All of the patients that were injected and subsequently went on to have a positive PCR test did not appear to experience severe symptoms as none were admitted to hospital. This study agrees with the national guidelines in that there is a risk of developing COVID-19 following a CSI however that risk in the studied population is low. Caution should be taken when considering CSI as a treatment option with a thorough risk/benefit analysis considered. Greater emphasis on shared decision making should be carried out with alternatives discussed prior to contemplating CSI to minimise the risk posed but there does not appear to be an increased risk of contracting COVID due to a CSI using Triamcinolone Acetonide for routine MSK conditions.

AUTHOR CONTRIBUTIONS

Daniel Lethbridge: Conceived and designed the analysis; Collected the data; Contributed data or analysis tools; Performed the analysis; Wrote the paper. **Simon O'Shea:** Conceived and designed the analysis; Contributed data or analysis tools; Other contribution.

CONFLICTS OF INTEREST

We confirm that this work is original and has not been published elsewhere nor is it currently under consideration for publication elsewhere. The authors declare that there are no conflicts of interest.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.

ETHICS STATEMENT

Hereby, I, Daniel Lethbridge consciously assure that for the manuscript Steroid injections in musculoskeletal conditions and COVID infection rates: What is the impact on positive rates following the injection? The following is fulfilled: (1) This material is the authors' own original work, which has not been previously published elsewhere. (2) The paper is not currently being considered for publication elsewhere. (3) The paper reflects the authors' own research and analysis in a truthful and complete manner. (4) The paper properly credits the meaningful contributions of co-authors and co-researchers. (5) The results are appropriately placed in the context of prior and existing research. (6) All sources used are properly disclosed (correct citation). Literally copying of text must be indicated as such by using quotation marks and giving proper reference. (7) All authors have been personally and actively involved in substantial work leading to the paper, and will take public responsibility for its content. The violation of the Ethical Statement rules may result in severe consequences. I agree with the above statements and declare that this submission is accurate.

ORCID

Daniel Lethbridge  <https://orcid.org/0000-0002-0525-5630>

REFERENCES

- Aziz, M. A., Hanifah, R. A., & Nahar, A. M. M. (2021). Musculoskeletal corticosteroid injection during COVID pandemic in Sahbah: Is it safe? *Advances in Orthopedics*, 2021, 1–6. <https://doi.org/10.1155/2021/8863210>
- Becker, D. E. (2013). Basic and clinical pharmacology of glucocorticosteroids. *Anesthesia Progress*, 60(1), 25–32. <https://doi.org/10.2344/0003-3006-60.1.25>
- British Pain Society. (2020). Pain management during COVID-19 viral infection. <https://www.britishpainsociety.org/static/uploads/resources/files/PainManagementduringCOVID-19viralinfection.pdf>
- British Society of Rheumatology. (2020). COVID-19 guidance. British Society of Rheumatology.
- British Society of Skeletal Radiology. (2020). The safety of corticosteroid injections during the COVID-19 global pandemic. <https://www.bssr.org.uk/static/uploads/forum/MusculoskeletalRadiologyduringtheCOVID-19GlobalPandemic.pdf3>
- Broersen, L. H., Pereira, A. M., Jørgensen, J. O., & Dekkers, O. M. (2015). Adrenal insufficiency in corticosteroids use: Systematic review and meta-analysis. *Journal of Clinical Endocrinology and Metabolism*, 100(6), 2171–2180. <https://doi.org/10.1210/jc.2015-1218>
- BSR BOA BASS RCGP BSIR FPM BPS CSP. (2020). Management of patients with musculoskeletal and rheumatic conditions who: Are on corticosteroids; require initiation of oral/IV corticosteroids; require a corticosteroid injection. [MSKcorticosteroidguidance.pdf \(londonhips.co.uk\)](https://www.londonhips.co.uk/MSKcorticosteroidguidance.pdf)
- Bugeja, M., Mariani, J., Dowling, J., Stringaro, G., Portelli, J. L., Sant, K., & Casaletto, J. (2021). Musculoskeletal steroid injections during the COVID-19 pandemic. *Journal of Orthopaedics*, 26, 103–106. <https://doi.org/10.1016/j.jor.2021.07.017>
- Cao, X. (2020). COVID-19: Immunopathology and its implications for therapy. *Nature Reviews Immunology*, 20(5), 269–270. <https://doi.org/10.1038/s41577-020-0308-3>
- Chang, C. Y., Prabhakar, A., Staffa, S. J., Hussein, J. S., Kheterpal, A. B., Simeone, F. J., & Bredella, M. A. (2021). Symptomatic COVID-19 infections in outpatient image-guided corticosteroid injection patients during the lockdown phase. *Skeletal Radiology*, 50(6), 1117–1123. <https://doi.org/10.1007/s00256-020-03656-w>
- Cruz-Topete, D., & Cidowski, J. A. (2015). One hormone, two actions: Anti- and pro-inflammatory effects of glucocorticoids. *Neuroimmunomodulation*, 22(1–2), 20–32. <https://doi.org/10.1159/000362724>
- Faculty of Pain Medicine of the Royal College of Anaesthetists. (2020). FPM response to concern related to the safety of steroids injected as part of pain procedures during the current COVID-19 virus pandemic. <https://fpm.ac.uk/sites/fpm/files/documents/2020-03/FPM-COVID-19-Steroid-Statement-2020.pdf2>
- Fascia, D., Dalili, D., Rennie, W., Rowbotham, E., Carne, A., & Robinson, P. (2020). Recommendations of the British Society of Skeletal Radiologists: The safety of corticosteroid injections during the COVID-19 global pandemic. British Society of Skeletal Radiologists. <https://doi.org/10.6084/m9.figshare.12662642>
- Finney, L. J., Glanville, N., Farne, H., Aniscenko, J., Fenwick, P., Kemp, S. V., Trujillo-Torralbo, M. B., Loo, S. L., Calderazzo, M. A., Wedzicha, J. A., Mallia, P., Bartlett, N. W., Johnston, S. L., & Singanayagam, A. (2021). Inhaled corticosteroids downregulate the SARS-CoV-2 receptor ACE2 in COPD through suppression of type I interferon. *The Journal of Allergy and Clinical Immunology*, 147(2), 510–519.e5. <https://doi.org/10.1016/j.jaci.2020.09.034>
- Habib, G. S. (2009). Systemic effects of intra-articular corticosteroids. *Clinical Rheumatology*, 28(7), 749–756. <https://doi.org/10.1007/s10067-009-1135-x>
- McKean, D., Chung, S. L., Fairhead, R., Bannister, O., Magliano, M., Papanikitas, J., Wong, N., & Hughes, R. (2020). Corticosteroid injections during the COVID-19 pandemic: Experience from a UK centre. *Bone & Joint Open*, 1(9), 605–611. <https://doi.org/10.1302/2633-1462.19.BJO-2020-0130.R1>
- Morgan, C., & Dattani, R. (2020). Should I use steroid injections to treat shoulder pain during the COVID-19 pandemic? *JSES International*, 4(4), 709–712. <https://doi.org/10.1016/j.jseint.2020.07.023>
- Newton, A. C., Jones, G., Jones, J., Norris, R., & Barabas, A. G. (2021). Intra-articular corticosteroid injections during the COVID-19 lockdown period: A service evaluation. *Musculoskeletal Care*, 19(2), 236–243. <https://doi.org/10.1002/msc.1530>
- NHS Digital. (2021). Risk criteria. digital.nhs.uk/coronavirus/shielded-patient-list/risk-criteria
- NHS England. (2020). COVID-19 therapy: Corticosteroids including dexamethasone and hydrocortisone. NHS England.
- Office of National Statistics (ONS). (2020). Coronavirus (COVID-19) infection survey pilot [Internet]. <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddisorders/bulletins/coronaviruscovid19infectionsurveyypilot/31july2020>

- RECOVERY Collaborative Group, Horby, P., Lim, W. S., Emberson, J. R., Maffham, M., Bell, J. L., Linsell, L., Staplin, N., Brightling, C., Ustianowski, A., Elmahi, E., Prudon, B., Green, C., Felton, T., Chadwick, D., Rege, K., Fegan, C., Chappell, L. C., Faust, S. N., Jaki, T., ... Landray, M. J. (2021). Dexamethasone in hospitalized patients with Covid-19. *The New England Journal of Medicine*, 384(8), 693–704. <https://doi.org/10.1056/NEJMoa2021436>
- Regan, P., Elkhailifa, S., & Barratt, P. (2021). The systemic immunosuppressive effects of peripheral corticosteroid injections: A narrative review of the evidence in the context of COVID-19. *Musculoskeletal Care*, 20(3), 1–11. <https://doi.org/10.1002/msc.1603>
- Stephens, M. B., Beutler, A. I., & O'Connor, F. G. (2008). Musculoskeletal injections: A review of the evidence. *American Family Physician*, 78(8), 971–976. [afp20081015p971.pdf\(aafp.org\)](https://doi.org/10.2539/afp20081015p971.pdf(aafp.org))
- Watson, J., Whiting, P. F., & Brush, J. E. (2020). Interpreting a COVID-19 test result. *BMJ*, 369, m1808. <https://doi.org/10.1136/bmj.m1808>
- Yanes-Lane, M., Winters, N., Fregonese, F., Bastos, M., Perlman-Arrow, S., Campbell, J. R., & Menzies, D. (2020). Proportion of asymptomatic infection among COVID-19 positive persons and their transmission potential: A systematic review and meta-analysis. *PLoS One*, 15(11), e0241536. <https://doi.org/10.1371/journal.pone.0241536>
- Yu, L. M., Bafadhel, M., Dorward, J., Hayward, G., Saville, B. R., Gbinigie, O., Van Hecke, O., Ogburn, E., Evans, P. H., Thomas, N., Patel, M. G., Richards, D., Berry, N., Detry, M. A., Saunders, C., Fitzgerald, M., Harris, V., Shanyinde, M., de Lusignan, S., ... PRINCIPLE Trial Collaborative Group. (2021). Inhaled budesonide for COVID-19 in people at high risk of complications in the community in the UK (PRINCIPLE): A randomised, controlled, open-label, adaptive platform trial. *Lancet (London, England)*, 398(10303), 843–855. [https://doi.org/10.1016/S0140-6736\(21\)01744-X](https://doi.org/10.1016/S0140-6736(21)01744-X)
- Yuki, K., Fujiogi, M., & Koutsogiannaki, S. (2020). COVID-19 pathophysiology: A review. *Clinical Immunology*, 215, 108427. <https://doi.org/10.1016/j.clim.2020.108427>

How to cite this article: Lethbridge, D., & O'Shea, S. (2022). Steroid injections in musculoskeletal conditions and COVID infection rates: What is the impact on positive rates following the injection? *Musculoskeletal Care*, 1–8. <https://doi.org/10.1002/msc.1707>